



VISION ENHANCEMENT GUIDE

# Simple Steps to Success

#### **IDEAL PATIENT CRITERIA FOR PRESBYOPES:**

Symptomatic presbyopes, current contact lens wearers or those motivated to wear contact lenses, normal cornea, astigmatism ≤1.00D

#### **See Naturally**

Follow these best practices when fitting NaturalVue® (etafilcon A) Enhanced Multifocal 1 Day™ contact lenses—now with the TripleTear® Lubrication System and Ultra-Tapered Edge designed for optimal fit and comfort.

#### **Starting** Power



Refract to 20/15



Check that the patient is not overplussed by using the binocular Red/Green (Duochrome test) 1 CLICK INTO GREEN



Use the best corrected spectacle refraction (BCSR) with the full cylinder component



Enter the full spectacle refraction into the NaturalVue® Multifocal QuickStart Calculator<sup>1</sup>

### Settle for 10 minutes

Let lenses settle 10 minutes & distance vision will begin to clear

#### **Evaluate & Enhance** - Distance First; then Near

Do NOT over-refract with the phoropter, flippers, loose trial lenses or trial frames. Make ON-EYE Diagnostic Lens Changes, if necessary. You are likely only 0.25D away from SUCCESS:

- To enhance DISTANCE: CHANGE THE DIAGNOSTIC LENS POWER on DOMINANT eye only by -0.25D (up to -0.50D) one lens change at a time and then in the NON-DOMINANT eye by -0.25D only if needed.
- To enhance NEAR: CHANGE THE DIAGNOSTIC LENS POWER on NON-DOMINANT eye by +0.25D (up to +0.50D) one lens change at a time and then in the DOMINANT eye by +0.25D only if needed.

#### NaturalVue® Enhanced Multifocal 1 Day™ Contact Lenses



- For presbyopes, a 0.25D lens change can greatly impact vision. Listen to the patient. If the patient says vision is not clear, **change the on-eye lens power**. Make a 0.25D change in the power, **one eye at a time**, starting with -0.25D on the dominant eye to refine distance. To refine near, once distance is optimized, start with +0.25D change on the non-dominant eye.
- Patients in monovision or center near multifocal designs may not have been using their central vision in both eyes for distance. It may take them a few days to fully adapt to this change. (Please see Stepwise Approach)

## For any questions or issues regarding fitting:

- ✓ Verify best corrected spectacle refraction used
- ✓ Use Red/Green (Duochrome) test to verify Rx is not overplussed
- ✓ Use the NaturalVue<sup>®</sup> Multifocal QuickStart Calculator
- ✓ Check that lenses have settled 10 minutes

Issue	Change the on-eye lens, one eye at a time
Distance vision not clear	<b>Dominant:</b> Change diagnostic lens -0.25 (-0.50 if needed) <b>Non-Dominant:</b> Change diagnostic lens -0.25 only if needed
Near vision not clear	Non-Dominant: Change diagnostic lens +0.25 (+0.50 if needed)  Dominant: +0.25 only if needed
Glare, Halos, vision seems "different"	<b>Dominant:</b> Change diagnostic lens -0.25 (-0.50 if needed)  If Center Near MF wearer or monovision, advise may take 2-3 days to adapt
Issues with driving at night	<b>Dominant:</b> Change diagnostic lens -0.25 (-0.50 if needed)
Vision is clear but feels strange, 3D effect, things jumping out at them	<b>Dominant:</b> Change diagnostic lens -0.25 (-0.50 if needed)  If Center Near MF wearer or monovision, advise may take 2-3 days to adapt
Have been adding minus and distance still not clear	Recheck that Best Corrected Spectacle Refraction is used.  Duochrome is one click into green and NaturalVue® QuickStart  Calculator is used
Vision fluctuating, going in and out, feels like have to stare at something to focus; Signs of over-minus	Non-Dominant: Change diagnostic lens +0.25 (+0.50 if needed) Dominant: +0.25 only if needed

For additional fitting support, kindly contact Professional Service Specialist alex.tay@oculuslens.com, or visit <a href="https://global.vtivision.com/practitioner">https://global.vtivision.com/practitioner</a> for more information

